

# Southwest Pediatric

A S S O C I A T E S

## Dosing for Commonly Used Medications

Weight (lb)	Acetaminophen (Tylenol)		Ibuprofen (Advil/Motrin)			Diphenhydramine (Benadryl)
	160 mg/5 ml	80 mg chew	100 mg/5 ml	50 mg/1.25 ml	100 mg chew	12.5 mg/5 ml
6-11	1.25 ml (1/4 tsp)	n/a	n/a	n/a	n/a	n/a
12-17	2.5 ml (1/2 tsp)	n/a	2.5 ml (1/2 tsp)	1.25 ml	n/a	2.5 ml (1/2 tsp)
18-23	3.75 ml (3/4 tsp)	1 1/2	3.75 ml (3/4 tsp)	1.875 ml	n/a	3.75 ml (3/4 tsp)
24-35	5 ml (1 tsp)	2	5 ml (1 tsp)	2.5 ml	1	5 ml (1 tsp)
36-47	7.5 ml (1 1/2 tsp)	3	7.5 ml (1 1/2 tsp)	n/a	1 1/2	7.5 ml (1 1/2 tsp)
48-59	10 ml (2 tsp)	4	10 ml (2 tsp)	n/a	2	10 ml (2 tsp)
60+	<i>See package instructions</i>					
	<i>Not recommended &lt; 2 months of age unless directed</i>		<i>Not recommended &lt; 6 months unless directed Make a note of the concentration.</i>			<i>Not recommended &lt; 2 years unless directed</i>

Kelly Jolet, MD | Vaishalee Patil, MD | Haydee Rimer, MD | Christine Trevino, DO | Nick Wagner, MD